

Cosmic Flows on 100 h^{-1} Mpc scales

Hume A. Feldman^{*,1}, Michael J. Hudson^{†,2} & Richard Watkins^{‡,3}

^{*}*Department of Physics & Astronomy, University of Kansas, Lawrence, KS 66045, USA.*

[†]*Department of Physics and Astronomy, University of Waterloo, Waterloo, ON N2L 3G1, Canada.*

[‡]*Department of Physics, Willamette University, Salem, OR 97301, USA.*

emails: ¹feldman@ku.edu; ²mjhudson@uwaterloo.ca; ³rwatkins@willamette.edu

To study galactic motions on the largest available scales, we require bulk flow moments whose window functions have as narrow a peak as possible and having as small an amplitude as possible outside the peak. Typically the moments found using the maximum likelihood estimate weights do not meet these criteria. We present a new method for calculating weights for moments that essentially allow us to "design" the moment's window function, subject, of course, to the distribution and uncertainties of the available data.

On scales that are small compared to the Hubble radius, galaxy motions are manifest in deviations from the idealized isotropic cosmological expansion

$$cz = H_0 r + \hat{\mathbf{r}} \cdot [\mathbf{v}(\mathbf{r}) - \mathbf{v}(0)] . \quad (1)$$

Redshift-distance samples, obtained from peculiar velocity surveys, allow us to determine the radial (line-of-sight) component of the peculiar velocity of each galaxy:

$$v(r) = \hat{\mathbf{r}} \cdot \mathbf{v}(\mathbf{r}) = cz - H_0 r . \quad (2)$$

Galaxies trace the large-scale linear velocity field $v(r)$ which is described by a Gaussian random field that is completely defined, in Fourier space, by its velocity power spectrum $P_v(k)$. The Fourier Transform of the line-of-sight velocity is

$$\hat{\mathbf{r}} \cdot \mathbf{v}(\mathbf{r}) = \frac{1}{(2\pi)^3} \int d^3\mathbf{k} \hat{\mathbf{r}} \cdot \hat{\mathbf{k}} v(\mathbf{k}) e^{i\mathbf{k} \cdot \mathbf{r}} , \quad (3)$$

which provides us with the velocity power spectrum

$$\langle v(\mathbf{k}) v^*(\mathbf{k}') \rangle = (2\pi)^3 P_v(k) \delta_D(\mathbf{k} - \mathbf{k}') , \quad (4)$$

where δ_D is the Dirac delta function. In linear theory, the velocity power spectrum is related to the density power spectrum

$$P_v(k) = \frac{H^2}{k^2} f^2(\Omega_{m,0} \Omega_\Lambda) P(k) , \quad (5)$$

where H is the Hubble constant and f is the rate of perturbation growth at the present. Ω_m and Ω_Λ are the matter and dark energy density parameters today. The power spectrum provides a complete statistical description of the linear peculiar velocity field.

For our power spectrum model, we follow Eisenstein and Hu (1998) in writing

$$P(k) \propto \sigma_8 \Omega_m^{0.6} k^n T^2(k/\Gamma) , \quad (6)$$

where T is the transfer function, Γ is the power spectrum "shape" parameter and as usual, we normalize the power spectrum using the parameter σ_8 , the amplitude of matter density perturbations on the scale of 8 h^{-1} Mpc.

To find the maximum likelihood parameters from peculiar velocity surveys we start with catalog of peculiar velocities galaxies, labeled by an index n , positions $\mathbf{r}_n$, estimates

of the line-of-sight peculiar velocities S_n and uncertainties σ_n . We assume that the observational errors are Gaussian distributed. We then model the velocity field as a uniform streaming motion, or bulk flow (BF), denoted by $\mathbf{U}$, about which there are random motions drawn from a Gaussian distribution with a 1-D velocity dispersion σ_* . Then the likelihood function for the BF components is

$$\mathcal{L}(U_i) = \prod_n \frac{1}{\sqrt{\sigma_n^2 + \sigma_*^2}} \exp \left(-\frac{1}{2} \frac{(S_n - \hat{r}_{n,i} U_i)^2}{\sigma_n^2 + \sigma_*^2} \right). \quad (7)$$

where i go from 1 to 3 to specify the BF components. While this has been done previously for maximum likelihood estimates (MLE) of the bulk flow moments, the advantage of these new moments is that they have been designed to be sensitive *only* to scales of order $100h^{-1}\text{Mpc}$. Thus we will be able to probe these scales without having to worry about the influence of smaller scales. Further, by isolating the very large scale motions we will be able to put stronger constraints on power spectrum parameters.

Maximum likelihood solution for BF moments is

$$U_i = A_{ij}^{-1} \sum_n \frac{\hat{r}_{n,j} S_n}{\sigma_n^2 + \sigma_*^2}, \quad (8)$$

where

$$A_{ij} = \sum_n \frac{\hat{r}_{n,i} \hat{r}_{n,j}}{\sigma_n^2 + \sigma_*^2}. \quad (9)$$

The measured peculiar velocity of galaxy n

$$S_n = \hat{r}_{n,i} v_i(\mathbf{r}_n) + \epsilon_n, \quad (10)$$

where ϵ_n is a Gaussian with zero mean and variance $\sigma_n^2 + \sigma_*^2$. The theoretical covariance matrix for the BF components

$$R_{ab} = R_{ab}^{(v)} + R_{ab}^{(\epsilon)}. \quad (11)$$

The first term is given as an integral over the matter fluctuation power spectrum, $P(k)$,

$$R_{ab}^{(v)} = \frac{\Omega_m^{1.2}}{2\pi^2} \int_0^\infty dk \mathcal{W}_{ab}^2(k) P(k), \quad (12)$$

where the angle-averaged tensor window function is

$$\mathcal{W}_{ab}^2(k) = \sum_{n,m} w_{a,n} w_{b,m} \int \frac{d^2 \hat{\mathbf{k}}}{4\pi} \left(\hat{\mathbf{r}}_n \cdot \hat{\mathbf{k}} \hat{\mathbf{r}}_m \cdot \hat{\mathbf{k}} \right) \exp(i\mathbf{k} \cdot (\mathbf{r}_n - \mathbf{r}_m)). \quad (13)$$

In order to do better than the MLE solutions, we redesign the window functions by requiring that they have narrow peaks and small amplitude outside the peak. We start with the BF MLE weights

$$w_{i,n} = A_{ij}^{-1} \sum_n \frac{\mathbf{x}_j \cdot \mathbf{r}_n}{\sigma_n^2 + \sigma_*^2}, \quad (14)$$

which depends on the spatial distribution and errors. Now consider an ideal survey with very large number of points, isotropic distribution and a Gaussian falloff $n(r) \propto \exp(-r^2/2R_I^2)$ where R_I is the depth of the survey.

The moments are specified by the weights

$$u_i = \sum_n w_{i,n} S_n \quad (15)$$




Fig. 1: The window functions of the bulk flow component for $R_I = 50h^{-1}\text{Mpc}$ for all the catalogs we considered. The thick (thin) lines are the window functions for the MV (MLE) bulk flow components. The x,y,z-component are dash-dot, short dash, long dash lines respectively. The Thick solid line is the ideal window function.




Fig. 2: The Γ likelihood function for the composite catalogs.

that minimize the variance $\langle (u_i - U_i)^2 \rangle$. We expand the variance

$$\langle (u_i - U_i)^2 \rangle = \sum_{n,m} w_{i,n} w_{i,m} \langle S_n S_m \rangle + \langle U_i^2 \rangle - 2 \sum_n w_{i,n} \langle U_i S_n \rangle, \quad (16)$$

since the measurement error included in S_n is uncorrelated with the bulk flow U_i . For BF moments we can impose the constraint that

$$\lim_{k \rightarrow 0} \mathcal{W}_{ii}^2(k) = \sum_{n,m} w_{i,n} w_{i,m} \int \frac{d^2 \hat{k}}{4\pi} (\hat{\mathbf{r}}_n \cdot \hat{\mathbf{k}} \hat{\mathbf{r}}_m \cdot \hat{\mathbf{k}}) = 1/3. \quad (17)$$

We now minimize this expression with respect to $w_{i,n}$ subject to the constraint which we enforce using a Lagrange multiplier to get

$$\sum_m (\langle S_n S_m \rangle + \lambda P_{nm}) w_{i,m} = \langle S_n U_i \rangle. \quad (18)$$

We solve this to get the minimum variance (MV) weights.

To apply this formalism, we used both cluster and galaxy peculiar velocity surveys. Specifically, we used the Willick cluster survey⁴, the SC1⁵ and SC2⁶ (Tully-Fisher clusters); The EFAR survey⁷ (a Fundamental Plane cluster survey); the SN are Type Ia supernovae from the compilation of Tonry *et al.*⁸; SMAC^{9,10} (a Fundamental Plane cluster survey); SFI++^{11,12} (a Tully-Fisher based survey of ~ 2700 field galaxies and ~ 730 groups).

The minimum variance, or MV, weights were calculated for the bulk flow component moments using the method described above for each of our catalogs, as well as for three composite catalogs, SFI++, consisting of both fields and groups from the catalog, OTHER, consisting of all the surveys except for the SFI++ field galaxy and groups catalogs, and Composite, consisting of all of the catalogs combined. The OTHER catalog was included due to the fact that, given their large size, the SFI++ catalogs tend to dominate any composite catalog in which they are included. Here we will show results from a deep survey, $R_I = 50 h^{-1}\text{Mpc}$.




Fig. 3: The bulk flow of the large catalogs. The triangles (points) are the BF moments with $R_I = 20h^{-1}\text{Mpc}$ ($R_I = 50h^{-1}\text{Mpc}$). There is a consistent and robust flow exhibited in all catalogs to the negative Galactic y direction (upper right panel) for the large scale analysis which is reflected in the BF magnitude (lower right panel)

In Figure 1 we show the window functions of the MV bulk flow component moments. For comparison, we also include the ideal window functions as well as those for the MLE moments for each survey. As expected, the match between the window functions for the MV moments and the ideal is best for the large surveys and those with small measurement error and similar distribution to the ideal survey. For the sparse, noisy surveys, the window functions for the MV moments are not very different than those of the MLE moments, differing mostly in the amplitude of the tail of the window function for large k .

In Figure 2. we show the likelihood function of the “shape” parameter Γ for the large catalogs. We see clearly that Γ is smaller for large R_I than for smaller R_I . In Figure 3. we show the BF velocity for the composite surveys. We see that for the shallow window functions, we get small flows, ~ 150 km/s. On large scales we get very large flows ~ 400 km/s. This unexpected flow which is exhibited in all the catalogs we analyzed, require a very steep power spectrum which leads to small Γ (see figure 2.)

References

1. Eisenstein D. J. & Hu, W., *ApJ* **496**, 605 (1998)
2. Giovanelli, R., *et al.* *AJ* **116**, 2632 (1998),
3. Tully, B. *et al.* *ApJ* **676**, 184 (2008).
4. Willick, J. A., *ApJ* **516**, 47 (1999)
5. Giovanelli, R. *et al.* *AJ* **116**, 2632 (1998)
6. Dale, D. A. *et al.* *AJ* **118**, 1489 (1999)
7. Colless, M. *et al.* *MNRAS* **321**, 277 (2001)
8. Tonry, J. L. *et al.* *ApJ* **594**, 1 (2003)
9. Hudson, M. J. *et al.* *ApJL* **512**, L79 (1999)

10. Hudson, M. J. *et al.* *MNRAS* **352**, 61 (2004)
11. Masters, K.L., *et al.* *ApJ* **653**, 861 (2006)
12. Springob, C. M. *et al.* *ApJS* **in press**, (2008)